

# Stopain® Clinical Gel - Product Information Sheet

Formulated with 10% menthol—plus dimethylsulfone (MSM) and glucosamine—**Stopain Clinical Gel** works on contact to provide fast, long-lasting pain relief of sprains, strains, bruises, arthritis, muscle aches, joint and back pain.

**Available in 3 industrial-sized applications for the professional setting:**

- 16 fl. oz. bottle
- 32 fl. oz. bottle
- 1 gallon (128 fl. oz.) bottle

**At-home sizes available for clients:**

- 4 fl. oz. tube
- (100) 5g packets in sample box



## BENEFITS

- Sweat-resistant
- Pharmaceutical grade skin conditioners
- Paraben-free
- No dyes or fragrances
- Non-greasy and Non-staining
- Pharmacist recommended

## ACTIVE INGREDIENTS

Menthol 10.0%

## PURPOSE

Topical  
Analgesic

## INACTIVE INGREDIENTS

Acrylic acid/vinyl ester copolymer, dimethylsulfone (MSM), eucalyptus oil, glucosamine sulfate, hydroxypropylcellulose, SD alcohol 39C, PEG-8 dimethicone, pentylene glycol, peppermint oil, triethanolamine, water (USP).

## DIRECTIONS

**Adults and children over 12 years:** apply directly onto affected area without the need to bandage. Repeat as necessary, but do not apply more than 4 times daily.

**Children 12 years or younger:** ask a doctor.

## WARNINGS

For external use only. Avoid contact with eyes. Flammable: Keep away from fire or flame.

**When using this product:**

- use only as directed
- do not bandage tightly or use a heating pad
- do not apply to wounds or damaged skin

**Stop use and ask a doctor if:**

- condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days
- redness is present
- excessive irritation of the skin develops

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

## OTHER INFORMATION

Store at room temperature.